

Effect of a High-Protein, Low-Carbohydrate Diet on Blood Glucose Control in People With Type 2 Diabetes

Mary C. Gannon^{1,2,3} and Frank Q. Nuttall^{1,3}

There has been interest in the effect of various types and amounts of dietary carbohydrates and proteins on blood glucose. On the basis of our previous data, we designed a high-protein/low-carbohydrate, weight-maintaining, nonketogenic diet. Its effect on glucose control in people with untreated type 2 diabetes was determined. We refer to this as a low-biologically-available-glucose (LoBAG) diet. Eight men were studied using a randomized 5-week crossover design with a 5-week washout period. The carbohydrate:protein:fat ratio of the control diet was 55:15:30. The test diet ratio was 20:30:50. Plasma and urinary β -hydroxybutyrate were similar on both diets. The mean 24-h integrated serum glucose at the end of the control and LoBAG diets was 198 and 126 mg/dl, respectively. The percentage of glycohemoglobin was 9.8 ± 0.5 and 7.6 ± 0.3 , respectively. It was still decreasing at the end of the LoBAG diet. Thus, the final calculated glycohemoglobin was estimated to be ~ 6.3 – 5.4% . Serum insulin was decreased, and plasma glucagon was increased. Serum cholesterol was unchanged. Thus, a LoBAG diet ingested for 5 weeks dramatically reduced the circulating glucose concentration in people with untreated type 2 diabetes. Potentially, this could be a patient-empowering way to ameliorate hyperglycemia without pharmacological intervention. The long-term effects of such a diet remain to be determined. *Diabetes* 53:2375–2382, 2004

Data obtained in our laboratory (1–3) as well as from others (reviewed in 4) indicate that glucose that is absorbed after the digestion of glucose-containing foods is largely responsible for the rise in the circulating glucose concentration after ingestion of mixed meals. Dietary proteins, fats, and absorbed fructose and galactose resulting from the digestion of sucrose and lactose, respectively, have little effect on blood glucose concentration.

We and others also have reported that even short-term starvation (hours) results in a dramatic decrease in the blood glucose concentration in people with type 2 diabetes

From the ¹Metabolic Research Laboratory and the Section of Endocrinology, Metabolism and Nutrition, Department of Veterans Affairs Medical Center, Minneapolis, Minnesota; the ²Department of Food Science and Nutrition, University of Minnesota, Minneapolis, Minnesota; and the ³Department of Medicine, University of Minnesota, Minneapolis, Minnesota.

Address correspondence and reprint requests to Mary C. Gannon, PhD, Metabolic Research Laboratory (111G), VA Medical Center, One Veterans Drive, Minneapolis, MN 55417. E-mail: ganno004@umn.edu.

Received for publication 5 February 2004 and accepted in revised form 2 June 2004.

LoBAG, low biologically available glucose; NEFA, nonesterified fatty acid; SDTU, special diagnostic and treatment unit.

© 2004 by the American Diabetes Association.

(5). This seems to be due largely to a rapid, progressive decrease in the rate of glycogenolysis (5,6). Hepatic glycogen stores in turn are dependent on the content of carbohydrate in the diet (6). Thus, a reduced-carbohydrate diet should result in a lower overnight fasting glucose concentration.

To test the hypothesis that a diet that is low in carbohydrate and particularly low in food-derived glucose could lower both the fasting and the postprandial blood glucose in people with type 2 diabetes, we designed a low-carbohydrate diet in which readily digestible starch-containing foods have been de-emphasized. However, the carbohydrate content is sufficient to prevent ketosis. This is in contrast to the low-carbohydrate diets being advocated for weight loss (7). We refer to this as a low-biologically-available-glucose (LoBAG) diet. In our study, we also attempted to ensure weight stability. The effect of 5 weeks of this diet on percentage glycohemoglobin and 24-h glucose, insulin, C-peptide, β -hydroxybutyrate, glucagon, triacylglycerol, and nonesterified fatty acid (NEFA) profiles in eight men with untreated type 2 diabetes is reported. Urea, creatinine, uric acid, and other data related to the metabolism of protein after ingestion of the LoBAG diet will be reported in a subsequent publication.

RESEARCH DESIGN AND METHODS

Men with mild, untreated type 2 diabetes were studied in a special diagnostic and treatment unit (SDTU; similar to a clinical research center). All participants met the National Diabetes Data Group criteria for the diagnosis of type 2 diabetes (8). Participant characteristics are given in Table 1. The study was approved by the Department of Veterans Affairs Medical Center and the University of Minnesota Committees on Human Subjects, and written informed consent was obtained from all participants. The participants did not have hematologic abnormalities, kidney disease, liver disease, macroalbuminuria (>300 mg/24 h), congestive heart failure, or untreated thyroid disease. Before the study, all participants were interviewed to determine their physical activity profile and food aversions and to explain the study process and commitment in detail. Participants confirmed that they had been weight stable for at least 3 months. They were instructed to maintain their current activity level throughout the study. Two weeks before beginning the study, the participants completed a 3-day food frequency questionnaire, with one of the days being a Saturday or a Sunday. This information was used to calculate the total food energy necessary to maintain body weight. None of the participants were being treated with oral hypoglycemic agents or insulin at the time of enrollment in the study. A 5-week randomized, crossover study design was used with a 5-week washout period between diets.

The control (15% protein) diet was designed according to the recommendations of the American Heart Association (9) and the U.S. Department of Agriculture (10,11). The diet consisted of 55% carbohydrate, with an emphasis on starch-containing foods, 15% protein, and 30% fat (10% monounsaturated, 10% polyunsaturated, and 10% saturated fatty acid). A second diet was designed to consist of 20% carbohydrate, 30% protein, and 50% fat. The saturated fatty acid content of the test diet was $\sim 10\%$ of total food energy; thus, the majority of the fat was mono- and polyunsaturated. This diet is referred to in the text as the LoBAG diet. The composition of the diets is given in Table 2.

TABLE 1
Patient characteristics

Patient	Age (years)	Height [in (cm)]	Weight [lb (kg)]	BMI (kg/m ²)	tGHb (%)	Duration of diabetes (months)	Concomitant diseases	Medications
1	69	74 (188)	221 (100)	27	8.7	60	Hypertension, dyslipidemia, coronary heart disease	Simvastatin, lisinopril, rabeprazole, ASA
2	72	69 (165)	239 (109)	35	10.0	12	Chronic obstructive pulmonary disease	Terazosin
3	51	68 (173)	181 (82)	27	8.6	12	None	ASA, naproxen
4	66	72 (183)	196 (89)	27	9.0	180	Hypertension	None
5	82	71 (180)	204 (93)	28	11.2	48	None	Lisinopril, ASA
6	56	72 (183)	267 (121)	35	10.1	24	Obesity, dyslipidemia	None
7	51	66 (168)	195 (89)	31	10.0	14	None	ASA, naproxen
8	59	67 (170)	233 (106)	36	9.4	19	Hypertension, obesity	Lisinopril
Mean	63.3	70 (176)	217 (99)	31	9.6	46		
Range	51–82	66–74 (168–188)	181–267 (82–121)	27–36	8.6–11.2	12–180		

ASA, acetylsalicylic acid.

Participants were randomized to begin the study with either the LoBAG or the control diet by a flip of a coin. Six participants started on the LoBAG diet, and five participants started on the control diet. Unfortunately, three of the participants who started on the control diet did not complete the study for personal reasons (death of spouse, move across country, chose not to finish). Therefore, the data are presented on eight participants who completed both arms of the study. Participants were admitted to the SDTU on the evening before the study. The next day, standardized meals that contained 55% carbohydrate, 30% fat, and 15% protein were given for breakfast, lunch, and dinner at 0800, 1200, and 1800. Participants were asked to remain in the SDTU during the study period with minimal activity.

On the second day in the SDTU, standardized meals again were given. This diet was similar for both baseline studies and is referred to as “control/pre” and “LoBAG/pre” diet in the figures, depending on which study diet followed the inpatient stay. In addition to the meals at 0800, 1200 and 1800, snacks were given at 1600 and 2100. Blood was obtained fasting at 0730, 0745, and 0800, every 15 min for the first hour after meals, every 30 min for the next 2 h, and then hourly until the next meal. Blood was drawn at a total of 46 time points. After this 24-h data accumulation period, the participants were sent home with all of the necessary food for the next 2–3 days as appropriate for the diet to which they were randomized.

Participants returned to the SDTU every 2–3 days to pick up food and meet with the study dietitian. At that time, they provided a urine specimen for analysis of creatinine and urea to determine dietary compliance. They also were weighed and had blood pressure, total glycohemoglobin (tGHb), and blood glucose measured. If their body weight decreased or increased on two successive occasions, then the total food energy of the meals was increased or decreased as appropriate to attempt to maintain weight stability throughout the study. In addition, participants were interviewed regarding dietary compliance, questions or concerns about the study, etc. At the end of the 5-week period, the participants again were admitted to the SDTU and blood was drawn as described above. At this time, the control or LoBAG meals (breakfast, lunch, dinner, and snacks) were given, as appropriate.

The plasma glucose concentration and β -hydroxybutyrate concentration

TABLE 2
Composition of diets

	Control	LoBAG
Energy (kcal)	2,825	2,825
Protein (g)	106 (15%)	210 (30%)
Carbohydrate (g)	388 (55%)	142 (20%)
Monosaccharides (g)	64	31
Disaccharides	50	16
Fat (g)	94 (30%)	158 (50%)
Monounsaturated (g)	29	62
Polyunsaturated (g)	24	35
Saturated (g)	33	30
Cholesterol (mg)	375	441
Dietary fiber (g)	24	36




FIG. 1. A: Mean body weight while on the control (○) or LoBAG (●) diet. B: Plasma β -hydroxybutyrate concentration after 5 weeks on the control (○) or LoBAG (●) diet.




FIG. 2. A: Mean plasma glucose concentration before (Δ) and after (\circ) 5 weeks on the control diet. *Side graph:* Net and total 24-h integrated glucose area response. Area response was not significantly different. **B:** Mean plasma glucose concentration before (\blacktriangle) and after (\bullet) 5 weeks on the LoBAG diet. *Side graph:* Net and total 24-h integrated glucose area response. *Both the net and the total area responses were significantly lower after the LoBAG diet ($P \leq 0.05$).

were determined by enzymic methods using an Analox analyzer with an O_2 electrode (Analox Instruments, London, U.K.). %tGhb was measured by boronate-affinity high-performance liquid chromatography (BioRad Variant; BioRad Labs, Hercules, CA). Serum immunoreactive insulin was measured using a standard double-antibody radioimmunoassay method using kits produced by Incstar (Stillwater, MN). Glucagon and C-peptide were measured by radioimmunoassay using kits from Linco Research (St. Louis, MO) and Diasorin (Stillwater, MN), respectively. NEFAs were measured enzymically using a kit manufactured by Wako Chemicals (Richmond, VA). Weight was determined in street clothes without shoes on a digital scale (Scalitrone, White Plains, NY). Blood pressure was measured using a Dinemap instrument (Critikon/Mediq, Pennsauken, NJ).

The net 24-h incremental area responses were calculated using the overnight fasting value as baseline. Total 24-h area responses were calculated using zero as the baseline. Both area calculations were done using a computer program based on the trapezoid rule. Statistics were determined using Student's t test for paired variates, with the Statview 512+ program (Brain Power, Calabasas, CA) for the Macintosh computer (Apple Computer, Cupertino, CA). $P < 0.05$ is the criterion for significance. Data are presented as the mean \pm SE.

RESULTS

The average body weight was 219 ± 10 lb (99 ± 4.5 kg) and 216 ± 10 lb (98 ± 4.5 kg) at the beginning of the control and LoBAG diets, respectively (Fig. 1A). At the end of the 5 weeks on the control diet, the average body weight was 215 ± 10 lb (98 ± 4.5 kg). After 5 weeks on the LoBAG diet, the average weight was 212 ± 9 lb (96 ± 4.1 kg). Thus, the average body weight decreased by 4 lb (1.8 kg) during the 5-week study period, regardless of diet.

Urine ketones were monitored twice weekly while participants were on the LoBAG diet. They were always zero to trace using nitroprusside impregnated Ketostix (Bayer, Elkhart, IN). Twenty-four-hour urine ketones were identical at the beginning and the end of the LoBAG diet (196 ± 8 and 196 ± 9 $\mu\text{mol/l}$, respectively). Before and after the




FIG. 3. A: Mean serum insulin concentration before (Δ) and after (\circ) 5 weeks on the control diet. *Side graph:* Net and total 24-h integrated insulin area response. Area response was not significantly different. **B:** Mean serum insulin concentration before (Δ) and after (\bullet) 5 weeks on the LoBAG diet. *Side graph:* Net and total 24-h integrated insulin area response. *Both the net and the total area responses were significantly lower after the LoBAG diet ($P \leq 0.05$).

control diet, they were 187 ± 7 and 203 ± 10 $\mu\text{mol/l}$, respectively.

The mean fasting β -hydroxybutyrate concentration was 225 ± 15 $\mu\text{mol/l}$ after 5 weeks on the control diet (Fig. 1B). After 5 weeks on the LoBAG diet, the mean fasting concentration was 236 ± 27 $\mu\text{mol/l}$. The 24-h profiles were similar when the participants ingested either diet.

The mean fasting glucose concentration before starting the control diet was 180 ± 10 mg/dl (10 ± 0.6 mmol/l; Fig. 2A). After 5 weeks on the control diet, the fasting glucose concentration was decreased to 159 ± 11 mg/dl (8.8 ± 0.6 mmol/l), but this was not significant ($P = 0.66$). Before starting the LoBAG diet, the mean fasting glucose concentration was 167 ± 13 mg/dl (9.3 ± 0.7 mmol/l), similar to that before starting the control diet ($P = 0.24$). After 5 weeks on the LoBAG diet, the fasting glucose concentra-

tion was significantly decreased to 119 ± 7 mg/dl (6.6 ± 0.4 mmol/l; $P < 0.003$; Fig. 2B).

The mean 24-h integrated net glucose area responses were similar precontrol, pre-LoBAG, and postcontrol (681 ± 174 , 731 ± 159 , and 730 ± 236 mg \cdot h \cdot dl $^{-1}$ [38 ± 9.7 , 41 ± 8.8 , and 41 ± 13.1 mmol \cdot h \cdot l $^{-1}$], respectively; Fig. 2 *side graphs, left bars*). After 5 weeks on the LoBAG diet, the net mean 24-h integrated glucose area response was decreased by 77% (165 ± 59 mg \cdot h \cdot dl $^{-1}$) (9.2 ± 3.3 mmol \cdot h \cdot l $^{-1}$; $P < 0.02$).

Total 24-h integrated glucose area responses also were similar precontrol, pre-LoBAG, and postcontrol ($4,998 \pm 337$, $4,746 \pm 301$, and $4,554 \pm 347$ mg \cdot h \cdot dl $^{-1}$ [278 ± 18.7 , 264 ± 16.7 , and 253 ± 19.3 mmol \cdot h \cdot l $^{-1}$], respectively; Fig. 2 *side graphs, right bars*). The total area response after 5 weeks on the LoBAG diet was decreased significantly




FIG. 4. Mean %tGhb response during the 5 weeks of the control (○) or LoBAG diet (●). *The tGhb on the test diet was significantly lower at weeks 3, 4, and 5 vs. the control diet ($P \leq 0.05$).

($3,023 \pm 160 \text{ mg} \cdot \text{h} \cdot \text{dl}^{-1}$ [$168 \pm 8.9 \text{ mmol} \cdot \text{h} \cdot \text{l}^{-1}$]; $P < 0.0004$ vs. the 5-week postcontrol and $P < 0.0001$ vs. pre-LoBAG). On the basis of these integrated areas, the mean glucose concentration over the 24-h periods of study was reduced from 198 to 126 mg/dl ($11\text{--}7 \text{ mmol/l}$) after 5 weeks on the LoBAG diet, a 36% decrease ($P < 0.0001$).

The mean fasting insulin concentrations before and after 5 weeks on both the control and the LoBAG diets were identical ($12 \pm 2 \text{ } \mu\text{U/ml}$ [$72 \pm 12 \text{ pmol/l}$]; Fig. 3). The mean 24-h integrated insulin area response above the fasting value was similar after the pre- and postcontrol diet and pre-LoBAG diet ($534 \pm 73 \text{ } \mu\text{U} \cdot \text{h}^{-1} \cdot \text{ml}^{-1}$ [$554 \pm 84 \text{ } \mu\text{U} \cdot \text{h}^{-1} \cdot \text{ml}^{-1}$]; and $530 \pm 81 \text{ } \mu\text{U} \cdot \text{h}^{-1} \cdot \text{ml}^{-1}$ [$3,024 \pm 438, 3,324 \pm 504, \text{ and } 3,180 \pm 486 \text{ pmol} \cdot \text{h} \cdot \text{l}^{-1}$], respectively; Fig. 3 *side graphs, left bars*). It was decreased at 5 weeks on the LoBAG diet ($318 \pm 39 \text{ } \mu\text{U} \cdot \text{h}^{-1} \cdot \text{ml}^{-1}$ [$1908 \pm 702 \text{ pmol} \cdot \text{h} \cdot \text{l}^{-1}$]). This was a decrease of 40% from the pre-LoBAG value ($P < 0.01$). The mean 24-h total integrated insulin area response decreased by 25%.

The mean fasting C-peptide concentration before and after the control diet was 0.86 ± 0.08 and $0.91 \pm 0.08 \text{ pg/ml}$. It was 0.81 ± 0.09 and 0.92 ± 0.08 before and after the LoBAG diet (data not shown). The 24-h time course response was similar to the insulin response. The net C-peptide area response was decreased by 34% after 5 weeks on the LoBAG diet. This was statistically significant ($P < 0.05$).

The mean %tGhb was essentially unchanged during the 5 weeks on the control diet (Fig. 4). A decrease in tGhb was present 1 week after the institution of the LoBAG diet and became significant by 3 weeks on the diet. At the end of the 5-week period, the %tGhb had decreased 22%, from 9.8 ± 0.5 to $7.6 \pm 0.3\%$ ($P < 0.0007$).

The mean fasting glucagon concentrations were similar before and after both the control and the LoBAG diets ($95 \pm 11, 91 \pm 8, 91 \pm 7, \text{ and } 94 \pm 7 \text{ pg/ml}$, respectively; Fig. 5). After 5 weeks on the LoBAG diet, the glucagon

response was similar to the control for the first hour after breakfast. Subsequently, the glucagon concentration was higher at every time point until 0700 the next morning, except for one time point after dinner. Both the net and the total glucagon area responses were significantly increased after the LoBAG diet ($P < 0.05$).

The mean fasting NEFA concentrations were $765 \pm 67, 654 \pm 59, 718 \pm 70, \text{ and } 593 \pm 50 \text{ } \mu\text{Eq/l}$, before and after the control and LoBAG diets, respectively (data not shown). These differences were not statistically significant ($P > 0.05$). The 24-h excursions were similar on the control and LoBAG prediet days. When the LoBAG diet was ingested, the fasting NEFA was lower and the increase after the lunch meal was attenuated, as was the decrease before dinner. The rise after dinner was more rapid and reached a higher concentration.

The mean 24-h integrated net NEFA area responses were $-5,323 \pm 1,187, -2,468 \pm 693, -4,525 \pm 1,660, \text{ and } 80 \pm 1,809 \text{ } \mu\text{Eq} \cdot \text{h} \cdot \text{l}^{-1}$ before and after the control and LoBAG diets, respectively. The small positive area response after the LoBAG diet was statistically significantly different compared with the response before the LoBAG diet ($P < 0.05$). Total areas were not statistically different from one another.

The mean fasting triacylglycerol concentrations were $264 \pm 36, 226 \pm 32, 246 \pm 27, \text{ and } 149 \pm 23 \text{ mg/dl}$ before and after the control and LoBAG diets, respectively (Fig. 6). The fasting triacylglycerol concentration was significantly lower after 5 weeks on the LoBAG diet ($P < 0.05$). After ingestion of either diet, the triacylglycerol concentration increased until $\sim 1200\text{--}1400$, decreased at $2000\text{--}2200$, increased slightly at ~ 2400 , and subsequently returned to the fasting value by 0800 the next morning.

The mean 24-h integrated net triacylglycerol area response was not significantly different between diets. However, the mean 24-h integrated total area response was significantly lower after 5 weeks on the LoBAG diet ($P < 0.05$).

The total cholesterol concentrations were $195 \pm 7, 184 \pm 17, 188 \pm 10, \text{ and } 177 \pm 8 \text{ mg/dl}$ before and after the control and the LoBAG diets, respectively. The LDL cholesterol concentrations were $105 \pm 9, 102 \pm 2, 105 \pm 7, \text{ and } 110 \pm 6 \text{ mg/dl}$ before and after the control and the LoBAG diets, respectively. The HDL cholesterol concentrations were $38 \pm 1, 37 \pm 2, 37 \pm 2, \text{ and } 36 \pm 2$ before and after the control and the LoBAG diets, respectively. These total, LDL, and HDL concentrations were not significantly different between diets or before and after each diet.

DISCUSSION

We previously reported that a diet in which the protein content was increased from 15 to 30% of total food energy, with a corresponding decrease in carbohydrate content, resulted in a moderate but highly statistically significant mean decrease in glycohemoglobin (8.1–7.3%) after 5 weeks on the diet. This was the consequence of smaller postmeal glucose increases. The fasting glucose concentration was unchanged (12).

In the present study, the diet contained the same 30% of food energy as protein. However, the carbohydrate content was further reduced from 40 to 20% of total food energy. The control diet in both studies is a diet that is




FIG. 5. *A*: Mean plasma glucagon concentration before (Δ) and after (\circ) 5 weeks on the control diet. *Side graph*: Net and total 24-h integrated glucagon area response. Area response was not significantly different. *B*: Mean plasma glucagon concentration before (Δ) and after (\bullet) 5 weeks on the LoBAG diet. *Side graph*: Net and total 24-h integrated glucagon area response. *The net and total area responses were significantly higher after the LoBAG diet ($P \leq 0.05$).

recommended for the general population as a means of reducing one's risk for coronary heart disease (9).

In the present study, the lower carbohydrate diet not only reduced the postmeal glucose concentration but also considerably reduced the overnight fasting glucose concentration. It is interesting that the 29% decrease observed in the present study is similar to the 34% decrease that we observed previously after a 36-h fast in people with type 2 diabetes (5). The overall result was a striking decrease in the 24-h integrated glucose concentration (Fig. 2). In addition, the percentage of glycohemoglobin concentration at the end of the 5-week study period was decreased from a mean of 9.8 to 7.6 (Fig. 4).

The study was designed to be 5 weeks in duration because 33 days had been reported to be the half-time for glycohemoglobin to reach a new steady state (13). If this is

the case, then the anticipated final percentage of glycohemoglobin would be ~ 5.4 (i.e., $2.2 \times 2 = 4.4$; $9.8 - 4.4 = 5.4\%$).

We previously determined that with the glycohemoglobin method that we use, each 1% glycohemoglobin represents ~ 20 mg/dl glucose integrated over a 24-h period (14). Using this information and the 24-h integrated glucose concentration observed at the end of the 5 weeks on the LoBAG diet, the estimated final percentage of glycohemoglobin would be 6.3%. Thus, the dietary modification that we refer to as the LoBAG diet has the potential for normalizing or nearly normalizing the blood glucose in people with mild to moderately severe type 2 diabetes. Nevertheless, these results should be considered to be merely a proof of concept. Only men were studied, and the diet was highly controlled and was of a relatively short




FIG. 6. A: Mean serum triacylglycerol concentration before (Δ) and after (\circ) 5 weeks on the control diet. Side graph: Net and total 24-h integrated triacylglycerol area response. Area response was not significantly different. B: Mean serum triacylglycerol concentration before (\blacktriangle) and after (\bullet) 5 weeks on the LoBAG diet. Side graph: Net and total 24-h integrated triacylglycerol area response. *The total area response was significantly lower after the LoBAG diet ($P \leq 0.05$).

duration. Thus, the generalization of these results will depend on additional longer-term studies in which both men and women and different age and ethnic groups are included and a greater variety of foods are used. In addition, even though we attempted to keep the participants' body weight stable, the participants lost a mean of ~ 4 lb while ingesting both diets.

The present data also suggest that a diet modification in which the protein and the fat content is increased will facilitate an improvement in glycohemoglobin by the various pharmacological agents used to treat diabetes. However, this also remains to be determined.

The decrease in postmeal glucose concentrations observed can easily be explained by the smaller amount of carbohydrate in the diet and thus the smaller amount of glucose absorbed after ingestion of the meals. The reason

for the greatly decreased fasting glucose concentration is uncertain but is likely to be the consequence of a reduced store of glycogen and thus a decrease in glycogenolysis rate (5,6,15). A priori, there is no reason to suspect that the LoBAG diet would result in a decreased rate of gluconeogenesis. Indeed, current evidence indicates that gluconeogenesis remains constant irrespective of the amount of carbohydrate in the diet (16) or of the gluconeogenic substrate supplied (15,17,18).

The LoBAG diet resulted in a decrease in 24-h integrated insulin concentration. In our previous study in which the protein content of the diet was increased from 15 to 30% of total food energy, the 24-h integrated insulin concentration was slightly increased when compared with the same control diet used in the present study (12). This was expected because dietary protein strongly stimulates insu-

lin secretion in people with type 2 diabetes (19). The decrease in integrated insulin concentration in the present study undoubtedly is due to the reduced food-derived glucose content of the diet. Dietary fat does not stimulate insulin secretion (20), or it facilitates a modest increase (4,21,22). Fructose (2,23) and galactose (24) ingestion also results in only a small increase in insulin concentration.

The serum total, LDL, and HDL cholesterol concentrations did not change significantly when the fat content of the diet was increased from 30 to 50% of total food energy. Most likely, this was because the saturated fatty acid content was kept at 10% of energy in both diets. The triglyceride concentration decreased as expected with a reduction in carbohydrate in the diet (25). A decrease in triglyceride might have been expected to increase HDL cholesterol (26); however, this has not been a consistent finding (27–29).

The glucagon area response increased 2.5-fold after the LoBAG diet. This increase is less than the fourfold increase that we observed in our previous study (12). However, the difference in fold increase is due, in part, to a difference in the response to the control diets. The net area response to the three 15% protein meals (control meals) was less in the previous study compared with the present study (139, 127, and 160 vs. 413, 293, and 349 $\text{pg} \cdot \text{h} \cdot \text{ml}^{-1}$, respectively). Nevertheless, the actual 24-h integrated glucagon response also was higher in the current study (893 vs. 525 $\text{pg} \cdot \text{h} \cdot \text{ml}^{-1}$).

In summary, a LoBAG diet can dramatically reduce the 24-h integrated glucose concentration and consequently the percentage of glycohemoglobin in people with type 2 diabetes. These positive results occur without a significant change in serum lipids, except for a significant decrease in triacylglycerol concentration.

ACKNOWLEDGMENTS

This study was supported by grants from the American Diabetes Association, the Minnesota Beef Council, and the Colorado and Nebraska Beef Councils.

The superb technical assistance of Kelly Jordan Schweim and Heidi Hoover is gratefully acknowledged. We thank the participants for volunteering for these studies, Brenda Tisdale and the staff of the SDTU and the Clinical Chemistry Laboratory for excellent technical expertise, Dr. Michael A. Kuskowski for advice on the statistical analysis and presentation of the data, and Ann Emery for excellent secretarial support.

REFERENCES

- Nuttall FQ, Gannon MC, Wald JL, Ahmed M: Plasma glucose and insulin profiles in normal subjects ingesting diets of varying carbohydrate, fat and protein content. *J Am Coll Nutr* 4:437–450, 1985
- Gannon MC, Nuttall FQ, Krezowski PA, Billington CJ, Parker S: The serum insulin and plasma glucose responses to milk and fruit products in type 2 (non-insulin-dependent) diabetic subjects. *Diabetologia* 29:784–791, 1986
- Krezowski PA, Nuttall FQ, Gannon MC, Billington CJ, Parker S: The insulin and glucose responses to various starch containing foods in type 2 diabetic subjects. *Diabetes Care* 10:205–212, 1987
- Nuttall FQ, Gannon MC: Carbohydrates and diabetes. In *American Diabetes Association Guide to Medical Nutrition Therapy for Diabetes*. Franz MJ, Bantle JP, Eds. Alexandria, American Diabetes Association, 1999, p. 85–106
- Gannon MC, Nuttall FQ, Lane JT, Fang S, Gupta V, Sandhofer C: Effect of 24 hours of starvation on plasma glucose and insulin concentrations in

- people with untreated non-insulin-dependent diabetes mellitus. *Metabolism* 45:492–497, 1996
- Nilsson LH, Furst P, Hultman E: Carbohydrate metabolism of the liver in normal man under varying dietary conditions. *Scand J Clin Lab Invest* 32:331–337, 1973
 - Atkins RC: *Dr. Atkins New Diet Revolution*. New York, Avon Books, 1998
 - Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 21 (Suppl. 1):S5–S19, 1998
 - American Heart Association: Dietary guidelines for healthy American adults: a statement for physicians and health professionals by the Nutrition Committee. *Circulation* 74:1465A–1468A, 1986
 - US Department of Agriculture: *The Food Guide Pyramid*. Washington, DC, U.S. Government Printing Office, 1992
 - US Department of Agriculture, US Department of Health and Human Services: Nutrition and Your Health: Dietary Guidelines for Americans. Washington, DC, U.S. Government Printing Office, 1995
 - Gannon MC, Nuttall FQ, Saeed A, Jordan K, Hoover H: An increase in dietary protein improved the blood glucose response in people with type 2 diabetes. *Am J Clin Nutr* 78:734–741, 2003
 - Rech ME: Observations on the decay of glycated hemoglobin HbA_{1c} in diabetic patients. *Exp Clin Endocrinol Diabetes* 104:102–105, 1996
 - Nuttall FQ: A comparison of percent total glycohemoglobin with percent HbA_{1c} in people with and without diabetes. *Diabetes Care* 21:1475–1480, 1998
 - Gannon MC, Nuttall JA, Damberg G, Gupta V, Nuttall FQ: Effect of protein ingestion on the glucose appearance rate in subjects with type 2 diabetes. *J Clin Endocrinol Metab* 86:1040–1047, 2001
 - Bisschop PH, Pereira Arias AM, Ackermans MT, Endert E, Pijl H, Kuipers F, Meijer AJ, Sauerwein HP, Romijn JA: The effects of carbohydrate variation in isocaloric diets on glycogenolysis and gluconeogenesis in healthy men. *J Clin Endocrinol Metab* 85:1963–1967, 2000
 - Jahoor F, Peters EJ, Wolfe RR: The relationship between gluconeogenic substrate supply and glucose production in humans. *Am J Physiol* 258:E288–E296, 1990
 - Jenssen T, Nurjhan N, Consoli A, Gerich JE: Failure of substrate-induced gluconeogenesis to increase overall glucose appearance in normal humans: demonstration of hepatic autoregulation without a change in plasma glucose concentration. *J Clin Invest* 86:489–497, 1990
 - Nuttall FQ, Mooradian AD, Gannon MC, Billington C, Krezowski P: Effect of protein ingestion on the glucose and insulin response to a standardized oral glucose load. *Diabetes Care* 7:465–470, 1984
 - May JM, Williams RH: The effect of endogenous gastric inhibitory polypeptide on glucose-induced insulin secretion in mild diabetes. *Diabetes* 27:849–855, 1978
 - Gannon MC, Ercan N, Westphal SA, Nuttall FQ: Effect of added fat on the plasma glucose and insulin response to ingested potato in individuals with NIDDM. *Diabetes Care* 16:874–880, 1993
 - Gannon MC, Nuttall FQ, Westphal SA, Seaquist ER: The effect of fat with carbohydrate on plasma glucose, insulin, C-peptide and triglycerides in normal male subjects. *J Am Coll Nutr* 12:36–41, 1993
 - Nuttall FQ, Gannon MC, Burmeister LA, Lane JT, Pyzdrowski KL: The metabolic response to various doses of fructose in type II diabetic subjects. *Metabolism* 41:510–517, 1992
 - Ercan N, Nuttall FQ, Gannon MC, Redmon JB, Sheridan KJ: Effects of glucose, galactose and lactose on the plasma glucose and insulin response in persons with non-insulin-dependent diabetes mellitus. *Metabolism* 42:1560–1567, 1993
 - Parks EJ, Hellerstein MK: Carbohydrate-induced hypertriglycerolemia: historical perspective and review of biological mechanisms. *Am J Clin Nutr* 71:412–433, 2000
 - Garg A, Bonanome A, Grundy SM, Zhang ZJ, Unger RH: Comparison of a high-carbohydrate diet with a high-monounsaturated-fat diet in patients with non-insulin dependent diabetes mellitus. *N Engl J Med* 319:829–834, 1988
 - Bonanome A, Visona A, Lusiani L, Beltramello G, Confortin L, Biffanti S, Sorgato F, Costa F, Pagnan A: Carbohydrate and lipid metabolism in patients with non-insulin-dependent diabetes mellitus: effects of a low-fat, high-carbohydrate diet vs a diet high in monounsaturated fatty acids. *Am J Clin Nutr* 54:586–590, 1991
 - Rasmussen OW, Thomsen C, Hansen KW, Vesterlund M, Winther E, Hermansen K: Effects on blood pressure, glucose, and lipid levels of a high-monounsaturated fat diet compared with a high-carbohydrate diet in NIDDM subjects. *Diabetes Care* 16:1565–1571, 1993
 - Garg A, Bantle JP, Henry RR, Coulston AM, Griver KA, Raatz SK, Brinkley L, Chen Y-D, Grundy SM, Huet BA, Reaven GM: Effects of varying carbohydrate content of diet in patients with non-insulin-dependent diabetes mellitus. *JAMA* 271:1421–1428, 1994